







### Medicina nasogastric feeding tubes

Medicina nasogastric feeding tubes have the physical characteristics required for feeding, flushing and delivering medication.

The worldwide introduction of the ENFit connector reduces the risk of enteral feeding tube misconnections and improves patient safety throughout hospitals and around the globe.







# Nasogastric feeding tubes and accessories

Medicina has a comprehensive range of nasogastric and nasojejunal tubes to feed a wide variety of patients from neonates to adults both in hospital and at home.

Within this range are tubes to feed all types of patients from 400g neonates to large adults, in both long and short-term designs. There are three types described in this booklet and sections on how to select the right size and length for a particular patient.

Other sections show how to pass a tube and check that it is in the correct position to feed. We have also put together a variety of accessories to help with feeding, aspiration, flushing and the delivery of medicines. We will also cover the care and maintenace of tubes and suggestions on what to do when anything goes wrong.




### Tube selection

It is important to choose the correct size of feeding tube in relation to the size of the patient and also in relation to the function required.

Feeding tubes tend to have narrower bores than aspiration tubes and smaller patients tend to have narrower tubes. The length of tube must also be sufficient to reach the desired site, without leaving too much external length on the face.

Caution: Neonates can be at risk from the use of longer tubes instead of an extention set, as the tube can become wrapped around the child's neck.



### Diameter

In adults, on standard feed, the most commonly used size is 8FR. If higher density feeds or feeds with fibre are used then a 12-14FR tube is normally selected. In children the standard size is 6FR with 8FR for thicker feeds or fibre feeds. In neonates 4-5FR is standard and 6FR for higher densities.

|          | Standard | High density |
|----------|----------|--------------|
| Patient  | feeds    | feeds        |
| Neonates | 4-5FR    | 6FR          |
| Children | 6FR      | 8FR          |
| Adults   | 8FR-10FR | 12FR-14FR    |

### Length

It is important to take a measurement of the approximate depth of a feeding tube placed into the stomach. Normally the distance from the ear to the nose and then from the episternum to the nose are added together.

For jejunal feeding add approximately 30cm more to the distance. It is also important to have sufficient external length to reach and connect to a feeding pump or to allow sufficient height for bolus feeding. Typical lengths used in neonates are 50cm, in children 50-75cm and in adults 75-85cm. An adult jejunal tube would be about 120-130cm.




### Tube characteristics

Tubes should be soft and have a smooth surface to ease their passage. The tube and guidewire should have an ENFit connector which fits ENFit syringes and administration sets. This is to reduce the risk of enteral tube feeding misconnections and improve patient safety. The tube should be marked every cm along the length to reference the

correct depth of the tube. The tip of the tube should be rounded and usually two feeding ports are better than one in case one becomes blocked. Bolus tips are sometimes made but they add to the diameter and make the tubes harder to pass. Weights are sometimes added to help passage through the pylorus into the small bowel.



### Key features



### Distance marks

All Medicina nasogastric and nasojejunal tubes have centimetre depth markings starting 5cm from the tip of the tube and stopping 5cm from the enteral connector. This makes it easy for the user to record the amount of tube placed inside the patient when it is first inserted. This can then be compared to the depth before every feed or medication to check that no movement has taken place.

Working alongside manufacturers, Medicina modified the printing technique to ensure crystal clear marking throughout the length of the tube.

### Guidewire

All Medicina long term feeding tubes have a self-lubricated non kink guidewire. Therefore no flushing is required to remove the guidewire before confirmation of tube placement.

### Y-Port NG tubes


Medicina have now introduced a "Yport" option for the long-term access for the medications and

### Radiopaque stripe

Medicina has made two major changes to the radiopaque stripe. The amount of Barium sulphate in the stripe has been doubled from the normal 20% to 40%. It is the amount of Barium sulphate that makes the tube visible under X-ray. The width of the radiopaque stripe has also been doubled. This makes it much easier to see the tubes on X-ray even in difficult patients.

Medicina has added a solid piece of polyurethane containing 40% Barium sulphate at the tip of longterm tubes. This will help identify the tip of the tube under X-ray. radiopaque tubing

short-term tubes



9.7ml

Many countries are now moving to the ENFit ISO Standard on all enteral feeding equipment



| nasogastric tube<br>priming volumes |                |                |       |
|-------------------------------------|----------------|----------------|-------|
| French                              | 50cm           | 80cm           | 120cr |
| 4FR                                 | 0.25ml         | 0 <b>.</b> 6ml | 0.9ml |
| 5FR                                 | 0 <b>.</b> 5ml | 0 <b>.</b> 8ml | 1.1ml |
| 6FR                                 | 0.7ml          | 1.1ml          | 1.6ml |
| 8FR                                 | 0.9ml          | 1.4ml          | 2.1ml |
| 10FR                                | 1.7ml          | 2.8ml          | 4.2ml |



Documentation of tube placement is very important. The label makes recording all the important details easier.

**Note:** labels are only available with selected feeding tubes.



All Medicina tubes have a colour coded strap which allows Clinical staff to see the diameter of the tube at a glance. The tubes also have the French Gauge and length printed at the top of the tube next to the male connector.

| 4FR  | red         |  |
|------|-------------|--|
| 5FR  | grey        |  |
| 6FR  | light green |  |
| 7FR  | pink        |  |
| 8FR  | light blue  |  |
| 9FR  | dark blue   |  |
| 10FR | black       |  |
| 12FR | white       |  |
| 14FR | dark green  |  |
| 16FR | orange      |  |
| 18FR | red         |  |
| 20FR | yellow      |  |
|      |             |  |



pH testing is the first line test for tube position. The Medicina pH indicator strip acts as its own colour chart, doing away with the need to have access to the box.

The pH strips are available separately in boxes of 100 or 200.







### LG4/120 LG4/125 LG4/160 LG5/40 LG6/120 LG6/160 LG8/40 LG8/50 LG8/90 LG8/120 LG8/160 LG10/120 LG10/160 LG12/120 LG16/120 LG18/120

### weighted tubes

LG6/50W 6FR x 50cm LG6/80W 6FR x 80cm LG8/50W 8FR x 50cm LG8/80W 8FR x 80cm LG10/80W 10FR x 80cm LG12/80W 12FR x 80cm

## Short-term polyurethane nasogastric feeding tubes

They are generally used in neonates and children but can be used in adults who require short-term enteral support. The tubes are used to feed, deliver medicines and aspirate stomach contents.

The tubes have an ENFit connector which fits ENFit syringes and the female ends of pump administration sets. All tubes have a radiopaque stripe to confirm position with X-ray (40% Barium sulphate compounded into the tube). The tubing itself is smooth and carries clear distance marks. The markings start 5cm from the tip of the tube and stop 5cm from the enteral connector to ensure an accurate reference point for tube position.

The popular sizes are also available in Tungsten weighted form which can be useful for feeding into the small bowel or where X-ray confirmation is difficult.

These tubes offer a comfortable solution to the nutritional support of infants and children.

#### Caution

In patients who are ventilated, the tubes should be changed every 14 days for adults and every 7 days for children to avoid chest infections or as clinically required.

### Features

- ENFit connection
- Fully radiopaque
- Crystal clear cm markings
- Two side eyes for feeding
- Rounded tip to aid insertion
- Colour coded strap



### Long-term fine bore feeding tubes

These tubes are used to feed adults and children. They are available from 6FR - 14FR diameters with a variety of lengths from 55cm - 130cm. Some popular sizes in 8FR and 10FR are also available weighted.

medical and care of the eldery units, paediatric units and also community

ends of pump administration sets. They also have an ENFit guidewire holder allowing aspiration to confirm correct position using enteral syringes without removing the guidewire.

The tubing itself is smooth and carries clear distance marks. The markings start 5cm from the tip of the tube and stop 5cm accurate reference point for tube position.

All tubes are fully radiopaque (40% Barium sulphate compounded into the clear stripe so that any change in aspirate colour or feed interaction with medicine can be detected without the need for tube removal.

polyurethane containing 40% Barium sulphate at the tip of all tubes. This will help to identify the tip of the tube under

X-ray if first method is unsuccessful.

Medicina have also introduced a 'Y-Port' feeding tube to the range. This allows easy access for medications and flushing while maintaining a closed system.


All Medicina long-term feeding tubes have a self-lubricated guidewire, therefore no flushing is required to remove the guidewire before confirmation of tube placement.

The nasojejunal tubes have a single open end feeding port that allows them to be passed over an endoscopically or radiologically placed guidewire should this be needed.

#### **Features**

- ENFit connection
- Fully radiopaque
- Self-lubricated non kink guidewire
- Crystal clear cm markings
- Two side eyes for feeding\*
- Rounded tip to aid insertion\*
- Optional 'Y-Port' NG tube
- Colour coded strap
- 40% Barium tip

\*On nasogastric tubes only.



#### Caution:

In patients who are ventilated, the tubes should be changed every 14 days for adults and every 7 days for children to avoid chest infections or as clinically required.

### nasogastric tubes

NGP8/75L NGP8/85L NGP8/120L

NGP6/95YC NGP8/85LW

**NJP8/130L** 8FR x 130cm

### weighted nasojejunal tubes





## Nasogastric feeding/aspiration tubes

This range of tubes is designed to aspirate stomach contents in adults and children. However they can be used for short-term feeding.

They are available from 6FR-20FR with lengths of 80cm - 100cm. Typical users would be critical care areas, theatres and surgical wards. They can also be used on an emergency crash trolley.

The tubes have an ENFit connector which fits enteral syringes and the female ends of pump administration sets. In addition they have a funnel or bladder type connector to fit suction equipment. The tubing itself is smooth and carries clear distance marks to reference tube position. All tubes have a radioopaque stripe to confirm position. To confirm position, use a pH strip or X-ray if first method is unsuccessful.


These tubes are softer and smoother than normal PVC Ryles tubes, they are easier to pass and more comfortable for the patients.

#### Caution:

In patients who are ventilated, the tubes should be changed every 14 days for adults and every 7 days for children to avoid chest infections or as clinically required.

#### Features

- ENFit & Funnel connector
- Fully radiopaque
- Crystal clear cm markings
- Rounded tip to aid insertion
- Four suction/ feeding holes
- Colour coded connector
- Reversible ENFit connector





These tubes are used to feed into the small bowel and simultaneously aspirate the stomach.


There are two versions of the tubes depending on the method of placement.


The aspiration lumen of these tubes may be used to feed the patient as they recover. Care must be taken changing the labels on the tubes.

The first, EF2L has its guidewire inside for the diagrams.


The second EF2EL, has an external the Wire" placement technique either radiologically or by using the guidewire diagrams.

### EF2L Endoscopic Placement






Using the endoscope place the tube into the small bowel.





2. Pass the endoscope into the stomach and catch the suture with the forceps.



To keep the tube in position, carefully and slowly remove the endoscope with a twisting motion having extended scope

### EF2L Surgical Placement






locate the bolus tip through the stomach wall.

EF2L




3. The feeding lumen can then be manually advanced into the jejumum.




4. Confirm the position of the tube by X-ray and remove guidewire.


### EF2EL Endoscopic Placement




1. Introduce endoscope into small bowel and insert guidewire down the centre.



2. Place guidewire into jejunum and remove endoscope.




Insert tube into nostril and thread over guidewire. Draw back tube and wire taking care not to dislodge wire.




Thread tube down into jejunum.
 Remove guidewire and check position by X-ray.


### EF2EL Radiological Placement



1. Place the guidewire into the jejunum under radiological control.



Insert tube into nostril and thread over guidewire. Draw back tube and wire taking care not to dislodge wire.



3. Thread tube down into jejunum. Remove guidewire and check position by X-ray.



### Passing NG tubes



1 Wash hands



2. Prepare equipment and explain procedure



3. Clear nostrils



4. Measure depth A



5. Measure depth B and then add A + B and note the total depth



6. Stretch tube



7. Lubricate tip and pass tube




8. Drink or swallow



9. Hold tube and aspirate



10. Confirm position



11. Remove guidewire





12. Fix tube onto face



3. Flush tube and check

### Position confirmation





MED50L

50ml Enteral

MED100L

100ml Entera

Drainage Ba

MED250L

250ml Enteral

MEDEOOL

500ml Enteral

Drainage Bag

MED1000L 1000ml Entera

Drainage Bag

NP01

NP02

N01

INU1 INO2 Aduit Nose Plaster Paediatric Nose Plaste 200 pH Indicator Strips

100 pH Indicator Strips

## Aspiration and drainage

When aspirating to confirm position, follow the guidelines on page 12 for pH confirmation.

### Enteral drainage Tube bags acce

Enteral drainage bags are available which connect to the tubes and allow collection of drained fluids. These bags have an antibacterial filter so that drained fluids can also be returned to the patient when

### Tube accessories

The pH strips are available separately in a box of 100 and 200 units. The plasters are also available separately in bags of 100. The plasters are hypoallergenic and can be used in neonates, children and adults



Tubes can be accessed directly with the Medicina range of enteral syringes.

Liquid medicines should be used where possible. Soluble tablets can be used if they are thoroughly dissolved. Never crush enteric coated medicines.

Medicines can be drawn up using the medicine straw. If glass ampoules are used then a medicine filter straw is also

Several bottle adapters are available to fit medicine bottles. A universal bottle adapter fits most bottles. There are seven other bottle adapters, which allow the tamper proof capping of the bottle in

### Filters

Medicines can be drawn up using the filter straw. These filters are recommended when any medicines are supplied in glass ampoules



A new medicines needle is available to access multiple use vials. It has an ENFit connector for use with the Medicina enteral syringes.

### Bottle adapters

A universal bottle adapter fits most bottles. In the case of medicines for children there are seven press-in tamper proof bottle adapters for the most common suspension bottles. These allow the bottle adapter to stay in the bottle and the tamper-proof cap can be used to close the bottle in between doses. The smallest fits bottles from 10-11mm and the largest fits bottles from 24-26mm in diameter

#### Caution


Medicina does not recommend the use of Cup Fill Technique for the drawing up of any medicine.

| BA02L |  |
|-------|--|
| BA03L |  |
| BA04L |  |
| BA05L |  |
| BA06L |  |
| BA07L |  |

BA07L Adapter is only available for use with Oral Tip/ENFit compatible syringes

| BA08L |  |
|-------|--|
|       |  |

| MF01L |  |
|-------|--|
| MN01L |  |
| MN02L |  |



### Delivering small volume medicines

It is advisable to use the medicine straw or bottle adapters for small volumes of medicines as the straw will occupy the dead space of the syringe in the same way as the male connector on the feeding tube. In this way medicines can be delivered accurately without considering the dead space. For very small doses (less than 0.5ml) the 1ml Low Dose Syringe should be used. It is important to flush tubes before and after medicines. Flushing before medicines prevents reactions with feeds. Flushing after medicines clears the priming volume of the tube and prevents reactions with feeds.


### Care and maintenance

Medicina nasogastric tubes can be used for the maximum period stated on the packaging but they should always be carefully maintained.

### External care

The external portion of the tube should be cleaned daily with mild soap and water and dried. Particular attention should be paid to the internal threads of the purple ENFit connector. This should be flushed thoroughly with enough water to remove any debris.


Dressings or any fixation materials should be changed daily.



### Internal care

The tube should be flushed every 4-6 hours when feeding and every 8 hours if the tube is not in use. In addition the tube should be flushed before and after medicines and before and after feeds, or in line with local hospital policies.

Sterile water should be used in infants less than six months old and any patients who may be imuno-compromised or jejunally fed. All others may use tap water. Typical volumes would be 20-30ml in adults, 10-20ml in children and 5-10ml in neonates. Care should be taken with fluid restricted patients. Larger syringes produce lower pressure so adults and children would normally use a 60ml and neonates a 20ml syringe or in line with local policies.



### Tube blockage

Tubes will only block if they are not well maintained by flushing or by delivering inappropriate forms of medications. Liquid medicines or soluble tablets should be used where possible. Any other tablets or forms of medicines should be dissolved and thoroughly diluted in line with pharmacy advice or manufacturer's instructions.

If a tube becomes blocked, make sure the external portion is not kinked. Using warm water attempt to flush the tube. If this fails, then fizzy water or pancreatic enzmes may be used in line with local policies. If tubes are flushed normally in line with recommended guidelines the tubes should not become blocked.

### Tube dislocation

If a new tube becomes dislodged by more than a few centimeters, then it should be removed and re-passed. Older tubes pose a risk of infection and should be replaced. Local practice and policies may vary from this. It is important never to re-introduce a guidewire into a tube which is still in the patient. The tube must be removed from the patient, the guidewire inserted into the tube then the tube can be re-passed.



### Product range

**Please note:** All products are available for purchase. Those <u>underlined</u> indicate products that are most commonly ordered/stocked through AMSL. For more information please contact AMSL Customer Service on (02) 9882 3666.

#### **ENFit syringes**



**LPE01LD** 1ml low dose enteral syringe (100)



**LPE25LD** 2.5ml low dose enteral syringe (100)




**LPE05** 5ml enteral syringe (100)




**LPE10** 10ml enteral syringe (100)



**LPE20** 20ml enteral syringe (80)



LPE30 30ml enteral syringe (80)



**LPE60** 60ml enteral syringe (60)



LPE100 100ml enteral syringe (30)

### ENFit reusable syringes



LHE01LD 1ml low dose reusable enteral syringe (100)



**LHE25LD** 2.5ml low dose reusable enteral syringe (100)



**LHE05** 5ml reusable enteral syringe (100)



**LHE10** 10ml reusable enteral syringe (100)



LHE20 20ml reusable enteral syringe (80)



**LHE60** 60ml reusable enteral syringe (60)

Oral Tip/ENFit Compatible Syringes




**OTO1LD** 1ml low dose enteral syringe (100)



**OT25LD** 2.5ml low dose enteral syringe (100)

#### ENFit extension sets




**EP015L** 15cm long short extension set (50)

**EP075L** 75cm long extension set (50)

**EP075LW** 75cm long wide bore extension set (50)

**EP150L** 150cm long extension set (50)



**EP015LYW** 15cm long wide bore Y-site short extension set (50)

**EP075LY** 75cm long Y-site extension set (50)

<u>EP075LYW</u> 75cm long wide bore Y-site short extension set (50)

**EP150LY** 150cm long Y-site extension set (50)



BA07L 10.5-11mm Bottle Adapter (50)

Only available for use with Oral Tip/ENFit compatible syringes.

### ENFit short-term polyurethane feeding tubes



12FR x 120cm (50)

14FR x 80cm (50)

16FR x 120cm (50)

18FR x 120cm (50)

6FR x 50cm (50)

6FR x 80cm (50)

8FR x 50cm (50)

8FR x 80cm (50)

10FR x 80cm (50)

12FR x 80cm (50)

LG12/120

LG14/80

LG16/120

LG18/120

Weighted LG6/50W

**LG6/80W** 


**LG8/50W** 

**LG8/80W** 

LG10/80W

LG12/80W

### ENFit long-term polyurethane feeding tubes



5FR x 55cm (25)


NGP5/55L

| Nul 3/33L   | 31 11 X 330111 (23) |
|-------------|---------------------|
| NGP6/55L    | 6FR x 55cm (25)     |
| NGP6/75L    | 6FR x 75cm (25)     |
| NGP6/85L    | 6FR x 85cm (25)     |
| NGP8/55L    | 8FR x 55cm (25)     |
| NGP8/75L    | 8FR x 75cm (25)     |
| NGP8/85L    | 8FR x 85cm (25)     |
| NGP8/120L   | 8FR x 120cm (25)    |
| NGP10/85L   | 10FR x 85cm (25)    |
| NGP10/120L  | 10FR x 120cm (25)   |
| NGP12/85L   | 12FR x 85cm (25)    |
| NGP12/100L  | 12FR x 100cm (25)   |
| NGP12/120L  | 12FR x 120cm (25)   |
| NGP14/85L   | 14FR x 85cm (25)    |
| NGP16/85L   | 16FR x 85cm (25)    |
| NGP6/95YC   | 6FR x 95cm (25)     |
| NGP8/95YC   | 8FR x 95cm (25)     |
| NGP10/95YC  | 10FR x 95cm (25)    |
| NGP12/95YC  | 12FR x 95cm (25)    |
| Weighted    |                     |
| NGP8/85LW   | 8FR x 85cm (25)     |
| NGP10/85LW  | 10FR x 85cm (25)    |
| NUL 10/03EM | 101117 000111 (20)  |
|             |                     |

### ENFit short-term polyurethane nasojejunal tubes


| 6FR x 120cm (10)  |
|-------------------|
| 8FR x 120cm (10)  |
| 8FR x 130cm (10)  |
| 10FR x 130cm (10) |
| 12FR x 130cm (10) |
| 14FR x 130cm (10) |
|                   |
| 10FR x 130cm (10) |
| 12FR x 130cm (10) |
| 14FR x 130cm (10) |
|                   |

### ENFit nasogastric feeding/aspiration tubes



| RT6/80L   | 6FR x 80cm (50)   |
|-----------|-------------------|
| RT8/80L   | 8FR x 80cm (50)   |
| RT10/100L | 10FR x 100cm (50) |
| RT12/100L | 12FR x 100cm (50) |
| RT14/100L | 14FR x 100cm (50) |
| RT16/100L | 16FR x 100cm (50) |
| RT18/100L | 18FR x 100cm (50) |
| RT20/100L | 20FR x 100cm (50) |
|           |                   |


### ENFit double lumen feeding tubes



**EF2L** Standard Nasojejunal Feed/Aspiration Tube 7/9FR x 130cm (5)

**EF2EL** Endoscopic Nasojejunal Feed/Aspiration Tube 7/9FR x 130cm (5)

### Enteral syringe pump




P1000 Enteral Syringe Pump (1)

#### Accessories



NP01 Adult nose plaster (100)

NP02 Paediatric nose plaster (100)



**IN02** pH indicator strips (100)

ENFit Gastrostomy extension set

ENFit Gastric Decompression systems




**EV01L** 250ml Gastric

Decompression System (30)

**EV02L** 500ml Gastric

Decompression System (30)

MGS02L Bolus or Gravity
Gastrostomy
Extension Set (30)





MED50L 50ml Enteral Drainage Bag (50)

MED100L 100ml Enteral Drainage Bag (50)

MED250L 250ml Enteral Drainage Bag (50)

MED500L 500ml Enteral Drainage Bag (50)

MED1000L 1000ml Enteral Drainage Bag (50)

### Gastrostomy accessories



GT01 Medium Gastrostomy Tidy (1)

75cm - 114cm

**GT02** Large Gastrostomy Tidy (1)

117cm - 152.2cm





### **ENFit Bottle adapters**



**BA01L** Stepped Bottle Adapter (50)

**=** 

BA02L 15-16.5mm Bottle Adapter (50)

BA03L 18-19.5mm Bottle Adapter (50)



BA04L 25.5-27mm Bottle Adapter (50)



BA05L 18.5-20.5mm Bottle Adapter (50) (Oramorph)



**BA06L** 20-21.5mm Bottle Adapter (50)



BA08L 16.5-18mm Bottle Adapter (50)

### **ENFit Transitional Adapters**



Transition Adapter for Reverse Luer Tube to ENFit Syringe (50)



Transition Adapter for ENFit Tube to Reverse Luer Syringe (50)



Tansition Adapter for Large Cone Tube to ENFit Syringe/Set (50)



Tansition Adapter for ENFit Tube to Large Cone Syringe (50)



Transition Adapter for ENFit Tube to Enlock Giving Set/Syringe (50)



Transition Adapter for ENFit Tube to Oral Tip Syringe/Set (50)



ML01L Transition Adapter for Oral Tip Tube to ENFit Syringe/Set (50)



**LK01** 

Transition Adapter for Funnel Tube to ENFit Syringe (50)

### ENFit syringe accessories



C005L Medicine Straw (50)




**C020L** 20cm Milk Straw (50)

C014L 14cm Milk Straw (50)

**<u>C010L</u>** 10cm Milk Straw (50)



MF01L Filter Straw (50)



SA03L 4-way Stopcock (50)



MN01L Enteral Blunt Fill Needle (100)



MN02L Enteral Blunt Fill Needle with Filter (100)



M158/1LD Male Caps (single) (50)

M158/8LD Male Caps 50 packs of 8 (400)

### Colostrum collector



TA10 Colostrum Collector (50)

## Medicina nasogastric feeding tubes

The ENFit ISO Standard range of nasogastric feeding tubes have been developed to deliver enteral feeds, medicines and flushes safely to neonatal, paediatric and adult patients.



For more information on the Medicina range, please contact us on **(02) 9882 3666** or at **amsl@amsl.com.au** 

amsl.com.au

